

AD-A093 585

NAVAL POSTGRADUATE SCHOOL MONTEREY CA F/G 8/1  
MICROPLANKTONIC ATP-BIOMASS AND GTP-PRODUCTIVITY ASSOCIATED WIT--ETC(U)  
SEP 80 S H BRONSINK

F/G 8/1

UNCLASSIFIED

NL

1. 118

END  
DATE  
FILMED  
2-81  
PTIC

AD A093585

2

# NAVAL POSTGRADUATE SCHOOL

Monterey, California



LEVEL II

DTIC  
SELECTED  
JAN 09 1981  
S F

M

## THESIS

C  
MICROPLANKTONIC ATP-BIOMASS AND  
GTP-PRODUCTIVITY ASSOCIATED WITH  
UPWELLING OFF PT. SUR, CALIFORNIA.

by

14 Sherman Hughes/Bronsink

11 September 1980 1271

Thesis Advisor:

E. D. Traganza

Approved for public release; distribution unlimited.

81 1 08 017

## UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A093-585	
4. TITLE (and Subtitle) Microplanktonic ATP-Biomass and GTP-Productivity Associated with Upwelling off Pt. Sur, California		5. TYPE OF REPORT & PERIOD COVERED Master's Thesis; September 1980
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Sherman Hughes Bronsink		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School ✓ Monterey, California		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California		12. REPORT DATE September 1980
		13. NUMBER OF PAGES 70
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Adenosine Triphosphate	Phosphate	Nutrients
Guanosine Triphosphate	Chlorophyll	Nitrate
Productivity	Temperature	
Biomass	Upwelling	
Fluxes	Satellites	
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>→ Microplanktonic ATP-biomass and GTP-productivity were studied in upwelling features off Pt. Sur, California. Biomass, determined by adenosine triphosphate (ATP) and chlorophyll <u>a</u>, was found to have a definite preferential location in the strong gradients of temperature and nutrients on the equatorward edge of the feature. Productivity, as sampled using the new method of guanosine triphosphate (GTP) sampling was found to be high throughout the</p>		

**SECURITY CLASSIFICATION OF THIS PAGE/When Data Entered**

The measurement of productivity and "growth potential" using GTP and GTP/ATP ratios was found to be a valuable tool in examining rapidly changing upwelling features. Preliminary results obtained using the GTP technique suggest a relationship between the flux of nutrients across ocean fronts and planktonic productivity associated with frontogenic coastal upwelling systems.

1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100  
 101  
 102  
 103  
 104  
 105  
 106  
 107  
 108  
 109  
 110  
 111  
 112  
 113  
 114  
 115  
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156  
 157  
 158  
 159  
 160  
 161  
 162  
 163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202  
 203  
 204  
 205  
 206  
 207  
 208  
 209  
 210  
 211  
 212  
 213  
 214  
 215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224  
 225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244  
 245  
 246  
 247  
 248  
 249  
 250  
 251  
 252  
 253  
 254  
 255  
 256  
 257  
 258  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276  
 277  
 278  
 279  
 280  
 281  
 282  
 283  
 284  
 285  
 286  
 287  
 288  
 289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298  
 299  
 300  
 301  
 302  
 303  
 304  
 305  
 306  
 307  
 308  
 309  
 310  
 311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319  
 320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368  
 369  
 370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396  
 397  
 398  
 399  
 400  
 401  
 402  
 403  
 404  
 405  
 406  
 407  
 408  
 409  
 410  
 411  
 412  
 413  
 414  
 415  
 416  
 417  
 418  
 419  
 420  
 421  
 422  
 423  
 424  
 425  
 426  
 427  
 428  
 429  
 430  
 431  
 432  
 433  
 434  
 435  
 436  
 437  
 438  
 439  
 440  
 441  
 442  
 443  
 444  
 445  
 446  
 447  
 448  
 449  
 450  
 451  
 452  
 453  
 454  
 455  
 456  
 457  
 458  
 459  
 460  
 461  
 462  
 463  
 464  
 465  
 466  
 467  
 468  
 469  
 470  
 471  
 472  
 473  
 474  
 475  
 476  
 477  
 478  
 479  
 480  
 481  
 482  
 483  
 484  
 485  
 486  
 487  
 488  
 489  
 490  
 491  
 492  
 493  
 494  
 495  
 496  
 497  
 498  
 499  
 500  
 501  
 502  
 503  
 504  
 505  
 506  
 507  
 508  
 509  
 510  
 511  
 512  
 513  
 514  
 515  
 516  
 517  
 518  
 519  
 520  
 521  
 522  
 523  
 524  
 525

Approved for public release; distribution unlimited.

Microplanktonic ATP-Biomass and GTP-Productivity  
Associated with Upwelling off Pt. Sur, California

by

Sherman Hughes Bronsink  
Lieutenant, United States Navy  
B.S., Western Washington State College, 1973  
M.S., University of West Florida, 1975

Submitted in partial fulfillment of the  
requirements for the degree of

MASTER OF SCIENCE IN OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL  
September 1980

Author

Sherman Hughes Bronsink

Approved by:

Eugene A. Traganza

Thesis Advisor

E. C. Haderlie

Second Reader

Robert H. Bourke Jr.

Chairman, Department of Oceanography

William M. Latta

Dean of Science and Engineering

## ABSTRACT

Microplanktonic ATP-biomass and GTP-productivity were studied in upwelling features off Pt. Sur, California. Biomass, determined by adenosine triphosphate (ATP) and chlorophyll-a, was found to have a definite preferential location in the strong gradients of temperature and nutrients on the equatorward edge of the feature.

Productivity, as sampled using the new method of guanosine triphosphate (GTP) sampling was found to be high throughout the thermo-nutrient gradient associated with an upwelling feature in an early stage of development.

The measurement of productivity and "growth potential" using GTP and GTP/ATP ratios was found to be a valuable tool in examining rapidly changing upwelling features. Preliminary results obtained using the GTP technique suggest a relationship between the flux of nutrients across ocean fronts and planktonic productivity associated with frontogenic coastal upwelling systems.

## TABLE OF CONTENTS

I.	INTRODUCTION-----	11
	A. HORIZONTAL NUTRIENT FLUXES-----	12
	B. VERTICAL NUTRIENT FLUXES-----	13
	C. FRONTAL NUTRIENT FLUXES-----	14
	D. ASSAYING ATP AND GTP-----	14
	E. GUANOSINE TRIPHOSPHATE-----	17
II.	METHODS-----	18
	A. CHLOROPHYLL <u>a</u> -----	18
	B. ATP AND $\Delta$ ATP-----	19
	C. $\Delta$ ATP VS. GTP CORRELATION EXPERIMENT-----	21
III.	RESULTS-----	23
	A. $\Delta$ ATP VS. GTP CORRELATION EXPERIMENT-----	23
	B. CRUISES-----	24
	1. September and November Cruises-----	24
	2. June Cruise-----	52
IV.	DISCUSSION-----	55
	A. $\Delta$ ATP VS. GTP CORRELATION EXPERIMENT-----	55
	B. THE JUNE CRUISE-----	55
V.	CONCLUSIONS-----	62
	BIBLIOGRAPHY-----	65
	INITIAL DISTRIBUTION LIST-----	68

## LIST OF TABLES

I.	$\Delta$ ATP as a Measure of GTP in a Binucleotide Mixture-----	22
II.	Ranges of ATP, $\Delta$ ATP, $\Delta$ ATP/ATP, and Chlorophyll <u>a</u> for the June 11-12, 1980 Cruise-----	53
III.	Correlation Coefficients between ATP, $\Delta$ ATP, $\Delta$ ATP, Chlorophyll <u>a</u> , and Temperature for the June 1980 Cruise-----	54



## LIST OF FIGURES

1. Kinetics of ATP, GTP, UTP, and CTP-dependent light emissions using crude luciferase preparations--	16
2. Bathymetry of the study area-----	25
3. Cruise track for the September 1979 cruise-----	26
4. Nitrate, phosphate, and sea surface temperature versus elapsed distance along the track of the September 27-28, 1980 cruise-----	27
5. ATP, chlorophyll <u>a</u> , and sea surface temperature versus elapsed distance along the track of the September 27-28, 1980 cruise-----	29
6. ATP distribution for the September 1979 cruise-----	31
7. Sea surface temperature for the September 1979 cruise-----	32
8. Nitrate distribution for the September 1979 cruise---	33
9. Phosphate distribution for the September 1979 cruise-----	34
10. Cruise track for the November 1979 cruise-----	35
11. Nitrate and sea surface temperature versus elapsed distance along the track of the November 29-30, 1979 cruise-----	36
12. ATP, chlorophyll <u>a</u> , and sea surface temperature versus elapsed distance along the track of the November 29-30, 1979 cruise-----	37
13. ATP distribution for the November 1979 cruise-----	38
14. Sea surface temperature for the November 1979 cruise-----	39
15. Nitrate distribution for the November 1979 cruise----	40
16. Cruise track for the June 1980 cruise-----	41

17.	Nitrate, phosphate, and sea surface temperature versus elapsed distance along the cruise track of the June 10-11, 1980 cruise-----	42
18.	ATP, chlorophyll <u>a</u> , and sea surface temperature versus elapsed distance along the track of the June 1980 cruise-----	43
19.	ATP, $\Delta$ ATP/ATP ratio, and sea surface tempera- ture versus elapsed distance along the track of the June 1980 cruise-----	44
20.	ATP distribution for the June 1980 cruise-----	45
21.	$\Delta$ ATP/ATP ratio distribution for the June 1980 cruise-----	46
22.	$\Delta$ ATP distribution for the June 1980 cruise-----	47
23.	Chlorophyll distribution for the June 1980 cruise-----	48
24.	Sea surface temperature for the June 1980 cruise-----	49
25.	Nitrate distribution for the June 1980 cruise-----	50
26.	Phosphate distribution for the June 1980 cruise-----	51

## LIST OF PHOTOGRAPHIC PLATES

1. TIROS-N Satellite IR Image of the California  
Coast, 9 June 1980----- 56

## ACKNOWLEDGEMENTS

This thesis is a result of ongoing research in chemical oceanography at the Naval Postgraduate School which is supported by the Office of Naval Research, Code 482, NORDA, NSTL Station, Bay St. Louis, Mississippi. I thank the sponsors and key individuals whose assistance made this task possible, including: Dr. Eugene D. Traganza, my thesis advisor and principal investigator; Captain W. W. Reynolds and the crew of the R/V ACANIA; Dana Austin and Bonita Hunter of the Naval Postgraduate School. Lastly, I would like to give special thanks to Walter E. Hanson, without whose assistance I would have been unable to complete this thesis.

## I. INTRODUCTION

This thesis examines the possibility that optimal growth conditions for microplankton occur near oceanic fronts as a result of horizontal nutrient fluxes. These fluxes may result from the sharp nutrient and thermal gradients that define such fronts. The research conducted here is part of a continuing study of the oceanic fronts and eddies formed by upwelling off Pt. Sur, California. During previous research [Traganza, Nestor, and McDonald, 1980a; Traganza, Conrad, and Breaker, 1980b], comparison of satellite thermal imagery to in situ biomass, nutrient, and thermal measurements showed large spatial variations in biomass concentrations. Microplanktonic blooms appeared greatest on the equatorward edge of the curl of "cyclonic upwelling systems" and adjacent to the sharp gradients in nutrients and temperature. There are several possible reasons for this distribution. The biomass may be advected into the area where it is found. It may be there initially, or it may appear as a preferential growth response to the newly formed front.

Preferential growth should be the result of optimal environmental conditions. Since large microplanktonic blooms (as measured by chlorophyll a and adenosine triphosphate (ATP)) appeared near sharp gradients of nutrients and temperature between upwelled, nutrient-rich, biochemically new

water and nutrient depleted, biochemically old, oceanic water, a nutrient and thermal flux may provide the necessary growth conditions. A horizontal, outward flow from the upwelling system may maintain the gradients ("a natural chemostat") and simultaneously provide a mechanism for biological seeding of fronts with the newly upwelled organisms [Tragana, et al., 1980b].

#### A. HORIZONTAL NUTRIENT FLUXES

Other studies have related phytoplankton production to horizontal and vertical nutrient gradients in Monterey Bay [Lasley, 1977] and to vertical fluxes of nutrients off Southern California [Eppley, Renger, Harrison, 1979].

Lasley [1977] computed the variations of phosphate, nitrate, and chlorophyll a using the distribution of variables equation [Sverdrup, et al., 1942]:

$$\frac{\partial C}{\partial t} = -u \frac{\partial C}{\partial x} - v \frac{\partial C}{\partial y} - w \frac{\partial C}{\partial z} + K \frac{\partial^2 C}{\partial x^2} + R \quad 1)$$

where C represents the concentration of the variable (at 5m in Lasley's case), t is time, u, v, and w are water velocity in the x, y, and z directions (x defined as parallel to the coast, y perpendicular to the coast, and z as depth), K is the lateral eddy diffusion coefficient. R is a nonconservative term which represents uptake or assimilation of the given nutrient by phytoplankton.

From a knowledge of the lateral eddy diffusivity coefficient for Monterey Bay, the surface circulation and the nutrient distributions within the bay, it was possible to compute the nutrient fluxes and compare them to the biomass. Measurements of the movement of isopleths of nutrients provided actual flux measurements for comparison. A study of a bay system has several advantages over the study of an exposed coastal upwelling feature interacting with a Western boundary current. The diffusivity coefficient (K) was already known for this area. Surface circulation and nutrient fluxes could be measured with relative ease, since the "feature" (the bay) was a fixed one. The peak in phytoplanktonic biomass (as measured by chlorophyll a) was found near Pt. Pinos where the gradients of nutrients (and presumably the fluxes) were greatest.

#### B. VERTICAL NUTRIENT FLUXES

Eppley [1979] postulated that carbon assimilation and therefore productivity could be directly related to vertical nitrate fluxes (dN/dz) following the equation:

$$\mu P(\text{production} \cdot \text{m}^{-3}) \sim \frac{K_z (\text{dN/dz})}{z_0} \quad 2)$$

where P is phytoplankton standing stock ( $\text{mg/m}^3$ ),  $\mu$  is its specific growth rate ( $\text{day}^{-1}$ ) at depth  $z_n$ , and depth  $z_0$  (m) is the depth at which the nitrate concentration has increased

to  $1 \mu\text{g} \cdot \text{atom} \cdot \text{liter}^{-1}$ . The depth  $z_0$  is arbitrary and could be chosen for another concentration if the coefficient  $K_z$  were also changed.


### C. FRONTAL NUTRIENT FLUXES

In this study, biomass (ATP and chlorophyll a), nutrients (nitrates and phosphates), and sea surface temperature were collected on a developing upwelling feature detected by satellite (IR) imagery of the Central California coast on June 10-11, 1980. Productivity and growth potential (instantaneous productivity per unit biomass) were examined using the new method of guanosine triphosphate sampling [Karl, 1978].

Linear plots following the ship's track and surface maps of these parameters were constructed to show frontal locations (temperature and nutrients), concentrations of biomass (based on ATP and chlorophyll a), areas of high productivity (based on high  $\Delta\text{ATP}$ ; see below), and areas of high growth potential (based on high  $\Delta\text{ATP}/\text{ATP}$ ; see below).

### D. ASSAYING ATP AND GTP

The luciferin-luciferase assay principle for ATP [Holm-Hansen and Booth, 1966; Holm-Hansen and Karl, 1976] has been widely used in recent years. It involves measuring the light emission from the reaction





[Holm-Hansen and Karl, 1978] where  $LH_2$  is reduced luciferin, FL is firefly luciferase,  $PP_1$  is inorganic pyrophosphate, P is product, and  $h\nu$  is light energy.

It is relatively easy to extract ATP from "particulates" in sea water with the appropriate solvent. Once extracted, ATP can be measured and used to estimate living carbon biomass by application of a conversion factor representing the mean C/ATP ratio of the microbial (bacteria, algae, and microzooplankton) biomass [Holm-Hansen, 1969]. Even with the right solvent and appropriate analytical procedure there are problems associated with this assay method. The conversion factor is somewhat species dependant, and it is dependent on the energy charge (physiological state), and variation of the total adenylate pool,  $AMP + ADP + ATP$  [Traganza, 1979]. Diel and other effects have also been noted [Hunter, 1979].

Another problem area concerns the analytical interference of non-adenine nucleotide triphosphates (NTPs) in the assay [Rasmussen and Nielsen, 1968]. When the assay is conducted using crude firefly luciferase and integrated light readings (60 seconds or more of light emissions), Guanosine triphosphate (GTP), uridine triphosphate (UTP), and cystosine triphosphate (CTP) all alter the light emission characteristics. The effect is most pronounced when these NTPs are at or near the concentration of ATP. Crystalline luciferase enzyme is ATP specific [McElroy and Green, 1956]. Peak readings (reading the light emission over the first six

seconds of the reaction instead of over 60 seconds) minimizes the interference, because ATP reacts much more rapidly with luciferin than other NTPs [Figure 1].




Figure 1. Kinetics of ATP, GTP, UTP, and CTP-dependent light emission using crude luciferase preparations. The concentration of each nucleotide was  $4 \times 10^{-8}$  M. [From Karl, 1978].

Measurements of ATP using crude extract and integrated readings result in an overestimation of the amount of ATP present because of the presence of other NTPs. This effect can be used to determine GTP levels which, in turn, may prove to be a measurement of productivity. Karl [1979] measured ATP in intertidal sediment extracts using both peak and integrated methods. The difference between these two values ( $\Delta\text{ATP} = \text{ATP (integrated)} - \text{ATP (peak)}$ ) showed a high correlation ( $r = .96$ ) to the measured amount of GTP present. The  $\Delta\text{ATP}$  values were poorly correlated to any

other parameter in the experiment (ATP, location the sample was from, etc.). From this Karl concluded that  $\Delta$ ATP is useful as a relative measure of GTP, especially in intertidal extracts.

#### E. GUANOSINE TRIPHOSPHATE

The ability to easily measure GTP is important to the study of mesoscale ocean fronts because of its relation to productivity. GTP is known to be essential in the biosynthesis of protein. Specifically, it is necessary in the initiation of synthesis, the aminoacyl tRNA binding, and the translocation processes [Watson, 1977]. Since protein synthesis is a major factor in primary productivity, a correlation between GTP and primary productivity is expected. Studies have shown a high linear correlation between cellular growth rate and GTP concentration (when normalized to the amount of ATP present, i.e., GTP/ATP) for bacteria [Karl, 1978], algae [Iwamura, et al., 1963], protozoa [Echetebeu and Plesner, 1977], and even fungi [Constantini et al., 1977]. The ratio of GTP to ATP may represent a "potential growth index" which would be invaluable because it would allow rapid assessment of productivity over large areas. This capability, combined with the rapid measurements of sea surface temperature (SST) and nutrients (nitrate and phosphate) may allow determination of the relation of productivity to fluxes in our rapidly developing systems.

## II. METHODS

Surface values of fluorescence (principally a measure of chlorophyll a), ATP, and  $\Delta$ ATP (as an index of relative GTP levels) were obtained off Pt. Sur, California during the development of a "cyclonic upwelling system" [Traganza et al., 1980b]. Temperature and nutrients (nitrate and phosphate) were obtained by Hanson [1980] and used here for interpretation of these results.

Nutrients (nitrate and phosphate) were sampled at two minute intervals using a Technicon Autoanalyzer. Sea surface temperature (SST) was recorded continuously from an intake located at approximately 2.5 meters depth. Enhanced TIROS-N satellite images were obtained from the National Environmental Satellite Service (NESS) at Redwood City, California.

### A. CHLOROPHYLL a

Fluorescence was measured continuously during the cruises using a Turner Model III fluorometer and converted to pigment concentration (principally chlorophyll a) as described by Lorenzen [1966]. To calibrate the system, discrete 275 ml seawater samples were taken in triplicate every hour. These were filtered through Whatman 4.5 cm diameter G/FC (pore size ca 0.45  $\mu$ m) glass fiber filters. The filters were folded, placed in polyethelene bags, frozen and analyzed within two

weeks, using the methods of Strickland and Parsons [1968]. Chlorophyll a concentrations were determined from the fluorescence record at two minute intervals to correspond to the nutrient sampling rate. To generate an organic carbon equivalent for comparison with organic carbon derived from ATP, "chlorophyll a" concentrations were converted to carbon units (mg/l) by using the average C/chlorophyll conversion factor of 100 as proposed by Holm-Hansen [1969]. Eppley [1977] has recommended a conversion factor of  $54 \pm 17$ . The two will be compared in the discussion section.

#### B. ATP AND $\Delta$ ATP

ATP and  $\Delta$ ATP samples were taken at approximately 10 minute intervals. Fifty ml samples of sea water were filtered through a 200  $\mu$ m nylon screen and then through a 2.5 cm (diameter) Reeve Angel 984-H (ca. 0.45  $\mu$ m pore size) glass filter. The filter was then placed in 100°C Trizma (Tris (Hydroxymethyl) aminomethane and hydrochloride) buffer (pH 7.7) to extract the ATP and other nucleotides. The buffer was then frozen for later analysis. The modified ATP method of Holm-Hansen and Karl [1976] was used in the ATP analysis.

SAI photometers, Models 1000 and 3000 (SAI, 4060 Sorrento Valley Boulevard, San Diego, California, 92121) were used to determine ATP concentrations from peak height measurements of the light emitted from the samples with an extract of

firefly luciferin and luciferase. Comparisons to standards (30, 15, 9, 6, and 3 ng/l) yielded ATP values in ng/l. These were converted to organic carbon biomass (mg/l) by using the average C/ATP conversion factor of 250 proposed by Holm-Hansen [1969] for the "microbial biomass" (bacteria, algae, and microzooplankton).

A relative measure of GTP present was computed using the " $\Delta$ ATP" method of Karl [1978]. ATP values were determined using both peak (6 second) and integrated (15 second delay, 60 second assay) methods on an SAI ATP photometer. A strip chart was fitted to the photometer to record peak readings which were converted to ATP concentrations by comparison of peak heights to the peak height of the standards. Integrated readings were given by the photometer's digital readout and converted to ATP concentrations by comparison to the digital readouts of the known standards. The difference between these estimates of the ATP concentration in a given sample ( $\Delta$ ATP) was used as a relative measure of GTP in the sample. According to Karl, 40 to 50% of the  $\Delta$ ATP value is GTP (personal communication). The  $\Delta$ ATP values were normalized to the ATP by dividing by the amount of ATP present ( $\Delta$ ATP/ATP). This ratio provides an index of "potential growth" or an estimate of the instantaneous productivity per unit biomass.

### C. $\Delta$ ATP GTP CORRELATION EXPERIMENT

As a test of the validity of using  $\Delta$ ATP as a measure of GTP on our instruments, a  $\Delta$ ATP vs. GTP correlation experiment was run using the SAI model 3000 photometer. Known concentrations of GTP and ATP were mixed in Trizma to represent varying percentages and absolute concentrations which may be found in sea water samples (Table I). "Peak" and "integrated" values were recorded on a strip chart interfaced to the photometer.

TABLE I

ΔATP AS A MEASURE OF GTP IN A BINUCLEOTIDE MIXTURE

GTP (ng/l)	ATP (ng/l)	ΔATP (CPM*)	% GTP (by wt)
.3	5.7	7	5
.6	5.4	8	10
.9	8.1	11	10
1.5	4.5	6	25
1.5	13.5	30	10
1.5	28.5	49	5
2.25	6.75	21	25
3.0	3.0	2	50
3.0	27.0	76	10
3.75	11.25	66	25
4.5	4.5	31	50
7.5	22.5	182	25
7.5	7.5	123	50
9.0	0	0	100
11.25	4.75	137	75
15.0	15.0	269	50
15.0	0.0	79	100
22.5	7.5	329	75
30.	0	286	100

\*CPM = Counts Per Minute, proportional to the amount of light emitted in the reaction. Values shown are averages of up to four readings taken at each concentration.



### III. RESULTS

#### A. $\Delta$ ATP VS. GTP CORRELATION EXPERIMENT

A linear regression analysis was performed on values of  $\Delta$ ATP vs. GTP using a WANG Series 700 Advanced Programming Calculator.

When all concentrations of ATP and GTP were considered (66 data points), a correlation coefficient or  $r$  of .87 was obtained. Adenosine diphosphate (ADP), derived from ATP, is a necessary catalyst for GTP reaction, viz;



with no ATP present initially, the only source of ADP is the crude extract, which limits the reaction. If the values with no ATP present are excluded (leaving 57 data points), an  $r$  value of .92 was obtained. If mixtures with a very low ATP content yielding a  $\Delta$ ATP of less than 10 counts are also excluded (leaving 36 data points), an  $r$  value of .94 was obtained. (In the analysis of the June 10-11, 1980 cruise, only three data points of 189 had a  $\Delta$ ATP of less than 10 counts.)

## B. CRUISES

Linear plots of nutrients, temperature and biomass following the cruise tracks were computer constructed by an IBM 360. A VERSATEC plotter was used to construct surface contour maps of the parameters in the upwelling features studied. The contour maps were hand smoothed to minimize the effects of advection (the feature moved during the 40 hours that samples were taken which introduced structure into contour maps which did not exist in reality) and computer generated anomalies which occurred at the margins of the data field (the outside edges of the feature were distorted by the boundary conditions set into the computer). These tract plots and surface maps are presented in Figures 2 to 26.

### 1. September and November Cruises

Surface maps and linear track maps from the September 27-28, 1979 and November 29-30, 1979 cruises are presented for background (Figures 3 to 15). The November 29-30, 1979 cruise best illustrates the biomass patchiness that was encountered at the equatorward edge of a feature. The ATP measured biomass shows several "cells" (Figure 6). Biomass is highest along the equatorward front of the feature where it reaches over 400 ng ATP/l (100 g C/l). Other, earlier cruises [Traganza, Conrad, and Greker, 1980], also showed the highest concentrations of biomass along the equatorward edge of the upwelled feature. Without information on productivity (or "growth potential" analogous to assimilation




Figure 2. Bathymetry of the study area.




Figure 3. Cruise track for the September 1979 cruise.




Figure 4. Nitrate, phosphate, and sea surface temperature versus elapsed distance along the track of the September 27-28, 1980 cruise.




Figure 4. Continued.




Figure 5. ATP, chlorophyll a, and sea surface temperature versus elapsed distance along the track of the September 27-28, 1980 cruise.




Figure 5. Continued.







Figure 6. ATP distribution for the September 1979 cruise.






Figure 8. Nitrate distribution for the September 1979 cruise.




Figure 9. Phosphate distribution for the September 1979 cruise.




Figure 10. Cruise track for the November 1979 cruise.  
S represents a hydro station number.




Figure 11. Nitrate and sea surface temperature versus elapsed distance along the track of the November 29-30, 1979 cruise.




Figure 12. ATP, chlorophyll *a*, and sea surface temperature versus elapsed distance along the track of the November 29-30, 1979 cruise.




Figure 13. ATP distribution for the November 1979 cruise.






Figure 14. Sea surface temperature for the November 1979 cruise.




Figure 15. Nitrate distribution for the November 1979 cruise.




Figure 16. Cruise track for the June 1980 cruise.




Figure 17. Nitrate, phosphate, and sea surface temperature versus elapsed distance along the track of the June 10-11, 1980 cruise.




Figure 18. ATP, chlorophyll a, and sea surface temperature versus elapsed distance along the track of the June 10-11, 1980 cruise.




Figure 19. ATP,  $\Delta\text{ATP}/\text{ATP}$  ratio, and sea surface temperature versus elapsed distance along the track of the June 1980 cruise.










Figure 22.  $\Delta$ ATP distribution for the June 1980 cruise.




Figure 23. Chlorophyll distribution for the June 1980 cruise.






Figure 25. Nitrate distribution for the June 1980 cruise.




Figure 26. Phosphate distribution for the June 1980 cruise.

number - see comments below)) and surface circulation, the reason for this patchiness could not be considered.

## 2. The June Cruise

On June 10-11, 1980, in addition to temperature, nutrient, and ATP data, "chlorophyll a" data was collected and, for the first time,  $\Delta$ ATP and  $\Delta$ ATP/ATP values were computed. The range of these values, which were computed at approximately 10 minute intervals, are given in Table II along with chlorophyll a and ATP. There were 189 ATP,  $\Delta$ ATP, and  $\Delta$ ATP/ATP values on the June cruise.

TABLE II  
RANGES OF ATP,  $\Delta$ ATP,  $\Delta$ ATP/ATP AND CHLOROPHYLL a  
FOR THE JUNE 11-12, 1980 CRUISE

	<u>high</u>	<u>low</u>	<u>mean</u>	<u>standard deviation</u>
ATP (ng/l)	1549	48	278	$\pm 195$
$\Delta$ ATP (ng/l)	358	.6	83	$\pm 30$
$\Delta$ ATP/ATP	.57	-.01	.30	$\pm .11$
Chlorophyll <u>a</u> (mg/m )	5.59	.17	1.11	$\pm .83$

Correlation coefficients for various parameters were computed and are summarized in Table III.

TABLE III

CORRELATION COEFFICIENTS BETWEEN ATP,  $\Delta$ ATP,  $\Delta$ ATP/ATP, CHLOROPHYLL<sub>a</sub>, AND TEMPERATURE FOR THE JUNE 1980 CRUISE

<u>Correlation of</u>	<u>Coefficient</u>
ATP to CHL	.536
ATP to $\Delta$ ATP/ATP	.249
ATP to $\Delta$ ATP	.851
$\Delta$ ATP/ATP to CHL	.35
$\Delta$ ATP/ATP to TEMP	-.249
$\Delta$ ATP/ATP to ATP	.613
$\Delta$ ATP to TEMP	-.218
$\Delta$ ATP to CHL	.559



#### IV. DISCUSSION

##### A. $\Delta$ ATP VS. GTP CORRELATION EXPERIMENT

The high correlation of  $\Delta$ ATP to GTP shows that in a binucleotid solution  $\Delta$ ATP may be taken as a measure of the relative amount of GTP present. In field samples there are other NTPs present (most importantly UTP) which enter into the reaction. Fortunately, for a given "system" (one with a relatively constant mix of microbial species) the percent of the  $\Delta$ ATP value which is GTP may be assumed to be constant [Karl, personal communication]. This constant, as mentioned previously, is on the order of 40 to 50% of the signal.

##### B. THE JUNE CRUISE

The June cruise shows a feature in what appears to be the "initiation phase" [Conrad, 1980]. The thermal and nutrient gradients are strong and well defined, and there are strong correlations between nitrate and phosphate ( $r = .96$ ), nitrate and temperature ( $r = -.96$ ), and phosphate to temperature ( $r = -.92$ ). These all point to "biochemically new" water [Traganza, et al., 1980] which has not had its initial conditions (temperature and nutrient concentrations) significantly altered by dynamical and biological processes. The satellite imagery (Plate 1) also shows a thermal pattern which has the appearance of an upwelling feature which is just forming and beginning to develop a cyclonic curl.



Plate 1. TIROS-N satellite IR image of the California coast, 9 June 1980.

ATP values are high (above 600 ng/l) in two areas (Fig. 20), the area of Monterey Bay off Pt. Pinos described by Lasley [1977] and discussed in the introduction, and in a narrow band along the seaward edge of the feature off Pt. Sur (stretching from about (5, -3) to (-3, 9) on the contour grid). This band closely follows the intermediate temperature band paralleling the peripheral margin of the satellite inferred feature and is in an area of water which is from 10.5 to 11.5 °Centigrade. This relation of the biomass to intermediate temperatures can also be seen in the linear plots (Figure 18). The maximum ATP concentrations are on the equatorward edge of the developing cyclonic curl (compare the SST figure and the ATP figure) which appears to be developing over the Sur Canyon. The nutrients are of intermediate concentration. Nitrates vary from 8 to 15  $\mu\text{M}$  and phosphates from 1.2 to 1.8  $\mu\text{M}$  in this area. The feature appears to be beginning a cyclonic spinup which would further concentrate and move this biomass along the equatorward edge of the feature.

Chlorophyll shows essentially the same distribution (Figure 23) with the exception of a large "cell" centered about (-1, 6) on the contour grid. Values here reach 3.5  $\text{mg}/\text{m}^3$ . There are several possible explanations for this anomaly. ATP is subject to several variations that chlorophyll a is not. It is energy charge dependent and shows more diel variation than chlorophyll a does [Hunter, 1979].

In addition, the difference may come from sampling technique. The continuous flow of water through the fluorometer ("chlorophyll a" determination) is not prefiltered because this produces a clogging problem. This could result in phytoplankton (i.e., chain formers) larger than 200 microns being excluded from the ATP samples which are prefiltered, but not from the chlorophyll samples.

$\Delta$ ATP (or "relative GTP") was highly correlated to ATP ( $r = .85$ ). Since they are not measures of the same parameter (GTP  $\sim$  instantaneous growth rate; ATP  $\sim$  living biomass), it can be concluded that all of the biomass is in a relatively uniform productive state.

Cellular alkaline phosphatase may interfere with these field measurements. Alkaline phosphatase acts by selectively consuming ATP and GTP. ATP is consumed at a more rapid rate than GTP, destroying evidence of the initial relationship [Karl, 1980]. The results in our field measurements appeared to be too consistent with temperature and nutrient data to have been effected by this problem. However, a separate test as described by Karl [1980] has been conducted on our stored field samples by Dana Austin at the Naval Postgraduate School. There are no indications of alkaline phosphatase interference in six samples selected from representative locations.

The  $\Delta$ ATP/ATP ratio or "growth potential" provides information on the productivity state of the biomass. The "growth potential" is analogous to assimilation number

(mg C uptake/mg Chl/hr). The standard deviation of the  $\Delta\text{ATP}/\text{ATP}$  ratio is only 37% of the mean value. This compares to a standard deviation of 70% of the mean for the ATP and 75% of the mean for chlorophyll. In other words, the "growth potential" varied very little and therefore the biomass adjacent to the fronts is in a relatively uniform productivity state. All of the biomass that is located adjacent to the nutrient fronts is present because of real growth. If the biomass were advected into the area, it should not remain at a high, uniform level. The only logical conclusion is that the biomass must be present as the result of real growth, i.e., presumably a preferential growth response to a horizontal nutrient flux across the sharp nutrient gradients.

The distribution of  $\Delta\text{ATP}/\text{ATP}$  is also of interest. The "growth potential" is high in the same area of Monterey Bay off Pt. Pinos described by Lasley [1977] and in the upwelling feature observed off Pt. Sur. In the feature, the "growth potential" is highest just seaward of the biomass peak. This suggests that the middle of the biomass peak is less active than the edges. This is not unexpected. Areas of high biomass are self-limiting in terms of growth, since they often encounter a growth limit such as light limiting or a nutrient shortage (although our data does not show a nutrient shortage). This also points to real growth in the system

(at least in this early stage) because the distribution is as expected for a growing system.

The relationship of the biomass and growth potential to the front suggests that the gradients are an important part of the productivity. As mentioned previously, other authors [Lasley, 1977; Eppley, et al., 1979] have related biomass to gradients by considering the biomass and productivity to flux relationships. Both of these authors were able to infer the circulation to compute the flux from the gradients, something that we have been unable to do at this time. The dynamic features we are studying do not lend themselves to the techniques they employed. Lasley used a fixed feature (Monterey Bay) which had known circulation (from current studies and drifters). Eppley worked with vertical fluxes and was able to calculate the circulation by applying continuity equations. If the surface circulation were better known in upwelled features as described here, the possibility is strong that a relationship between nutrient fluxes and the productivity would be evident. The results would probably be similar to Lasley's, if the eddy diffusion term and the advection term parallel to the nutrient contours are negligible. This leaves a distribution of variables equation of:

$$\frac{\partial C}{\partial t} = -u \frac{\partial C}{\partial x} - w \frac{\partial C}{\partial z} + R \quad 6)$$

This equation relates the changes in concentration of the

parameter (nutrients in this case) directly to the fluxes associated with the vertical and horizontal gradients (with the exception, of course, of the biological term, R). If this equation is accepted as representing the behavior of the nutrients in the feature, there is a good qualitative agreement between the location of the biomass and productivity vs. the fluxes.

## V. CONCLUSIONS

1. The study of  $\Delta\text{ATP}$  and  $\Delta\text{ATP}/\text{ATP}$  ratios can provide valuable insight into productivity and growth potential. This method is especially valuable in studying dynamic features such as fronts and eddies, where a rapidly changing mesoscale environment and the need for large numbers of samples make more traditional methods unworkable.

2. Several conclusions about the initial or early phase of upwelling features can be drawn from the study of the June cruise.

a. Areas of high biomass are areas of high productivity. The biomass present appears to be there as the result of optimal growth conditions, not advection.

b. The highest growth is in the area surrounding the highest biomass, i.e., intermediate concentrations of biomass appear to be optimal. Even at this early stage in the feature, biomass is beginning to be limited in its growth by some factor.

c. Since both high biomass and high growth potential are found in the area of large nutrient and temperature gradients, the gradients must have a favorable impact on growth. A nutrient flux, as postulated earlier, may serve to connect the two. Unfortunately, until the surface circulation is better known, computation of surface fluxes is not possible.



If equation 6) is applicable to the June feature, however, the good correlation between productivity and fluxes is suggested qualitatively in the data if the velocities across the gradients are relatively constant.

d. With the exception of one chlorophyll a "cell" described earlier, the surface maps of ATP and chlorophyll a are essentially the same. A comparison of the surface maps and mean values suggests that the standard conversion ratios of C:ATP (250:1) and C:Chl (100:1) are not entirely accurate in this feature. If a C:Chl ratio of 60:1 were used instead of 100:1, the values would be in closer agreement, suggesting that the populations are largely phytoplankton. Eppley [1977] found the ratio of  $54 \pm 17$  for C:Chl in phytoplankton populations off Southern California which supports this suggestion. Without specific knowledge of C:Chl and C:ATP ratios in the populations under investigation, one cannot conclude that microzooplankton or bacterial populations are indicated rather than phytoplankton when ATP is larger than chlorophyll.

e. Satellite imagery may be used to infer potential biomass location and growth potential, at least in an initial stage of a feature's development but not all features may support biological activity. Studies of features in later stages of development may provide insight into the biomass and growth potential of older features. Applications of this technology may lead to the ability to remotely predict

productive regions for fisheries and other applications. Such predictions would be dependent on the source water depth (a function of the feature's age, the season, and the recent wind history [Hanson, 1980]).

## BIBLIOGRAPHY

- Conrad, J.W., Relationships Between the Sea Surface Temperature and Nutrients in Satellite Detected Oceanic Fronts, M.S. Thesis, Naval Postgraduate School, 1980.
- Constantini, M.G., Zippel, R., and Sturani, E., Biochimica et Biophysica Acta, V. 476, p. 272-278, 1977.
- Echetebe, C.O., and Plesner, P., "The Pool of Ribonucleotide Triphosphates in Synchronized *Tetrahymena pyriformis*," Journal of General Microbiology, V. 103, p. 389-392, 1977.
- Eppley, R.W., Harrison, W.G., Chisholm, S.W., and Stewart, E., "particulate Organic Matter in Surface Waters of Southern California and Its Relationship to Phytoplankton," Journal of Marine Resources, V. 35, p. 671-696, 1977.
- Eppley, R.W., Renger, E.H., and Harrison, W.G., "Nitrate and Phytoplankton Production in Southern California Coastal Waters," Limnology and Oceanography, V. 24, p. 483-494, 1979.
- Hanson, W.E., Nutrient Study of Mesoscale Thermal Features off Pt. Sur, California, M.S. Thesis, Naval Postgraduate School, 1980.
- Holm-Hansen, O. and Boothe, C.R., "The Measurement of ATP in the Ocean and Its Ecological Significance," Limnology and Oceanography, V. 11, p. 510-519, 1966.
- Holm-Hansen, O. and Karl, D.M., "Biomass and Adenylate Energy Charge Determination in Microbial Cell Extracts and Environmental Samples," Methods in Enzymology, V. LVII, p. 73-85, 1978.
- Holm-Hansen, O. and Karl, D.M., "Effects of Luciferin Concentration on Quantitative Assay of ATP Using Crude Luciferase Preparations," Analytical Biochemistry, V. 75, p. 100-112, 1976.
- Holm-Hansen, O., "Determination of Microbial Biomass in Ocean Profiles," Limnology and Oceanography, V. 14, p. 740-747, 1969.

Hunter, B.L., ATP/Carbon Ratios as a Measure of Phytoplankton Biomass and Productivity, M.S. Thesis, University of Hawaii, 1979.

Iwamura, T., Kanuzawa, T., and Kunazawa, K., Studies on Microalgae and Photosynthetic Bacteria, Japanese Society of Plant Psysiologists, University of Tokyo Press, p. 587-598, 1963.

Karl, D.M., "Occurrence and Ecological Significance of GTP in the Ocean and in Microbial Cells," Applied and Environmental Microbiology, p. 349-355, 1978.

Karl, D.M., "Adenosine Triphosphate and Guanosine Triphosphate Determinations in Intertidal Sediments," Methodology for Biomass Determinations and Microbial Activities in Sediments, ASTM STP 673, C.D. Littlefield and P.L. Seyfried, Eds., American Society for Testing and Materials, p. 5-20, 1979.

Karl, D.M., and Craven, D.B., "Effects of Alkaline Phosphatase Activity on Nucleotide Measurements in Aquatic Microbial Communities," Applied and Environmental Microbiology, V. 40, No. 3, September 1980.

Lasley, S.R., Nearshore Circulation and Aging of Water in Monterey Bay, M.A. Thesis, San Jose State University, 1977.

Lorenzen, C.J., "A Method for the Continuous Measurement of In Vivo Chlorophyll Concentration," Deep Sea Research, V. 13, p. 223-227, 1966.

McElroy, W.D., and Green, A., Archives for Biochemistry and Biophysics, V. 64, p. 257-271, 1956.

Rasmussen, H. and Nielsen, R., Acta Chemica Scandinavia, V. 22, p. 1745-1762, 1968.

Strickland, J.D.H., and Parsons, T.R., A Practical Handbook of Seawater Analysis, J.C. Stevenson (Ed.), Queen's Printer and Comptroller of Stationary, p. 49-62 and 71-76, 1968.

Sverdrup, H.U., Johnson, M.W., and Fleming, R.H., The Oceans, Their Physics, Chemistry, and General Biology, Prentice Hall, 1942.

Traganza, E.D., Naval Postgraduate School Technical Report No. 68-79-006, "Equilibrium Calculations of ATP as a Theoretical Basis for the 'Normal' Range of C/ATP Ratios in Marine Zooplankton," September, 1979.

Traganza, E.D., Nestor, D.A., and McDonald, A.K., "Satellite Observations of a Nutrient Upwelling off the Coast of California," Journal of Geophysical Research, V. 85 No. C7, p. 4101-4106, July 1980.

Traganza, E.D., Conrad, J.W., Breaker, L.C., "Satellite Observations of a 'Cyclonic Upwelling System' and 'Giant Plume' in the California Current," Coastal Upwelling, American Geophysical Union (in press), 1980.

Watson, J.D., Molecular Biology of the Gene, W.A. Benjamin, Inc., p. 331-338, 1977.

# INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, VA 22314	2
2. Library, Code 0142 Naval Postgraduate School Monterey, CA 93940	2
3. Chairman, Code 68 Mr Department of Oceanography Naval Postgraduate School Monterey, CA 93940	1
4. Chairman, Code 63 Ha Department of Meteorology Naval Postgraduate School Monterey, CA 93940	1
5. Dr. E. D. Traganza, Code 68 Tg Department of Oceanography Naval Postgraduate School Monterey, CA 93940	5
6. LT. S. H. Bronsink Helsuppron Three NAS North Island San Diego, CA 92135	3
7. Director Naval Oceanography Division Navy Observatory 34th & Massachusetts Avenue NW Washington, D.C. 20390	1
8. Commander Naval Oceanography Command NSTL Station Bay St. Louis, MS 39529	1
9. Commanding Officer Naval Oceanographic Office NSTL Station Bay St. Louis, MS 39529	1

- |     |  |   |
|-----|--|---|
| 10. | Commanding Officer<br>Fleet Numerical Oceanography Center<br>Monterey, CA 93940  | 1 |
| 11. | Commanding Officer<br>Naval Ocean Research & Development<br>Activity<br>NSTL Station<br>Bay St. Louis, MS 39529                            | 1 |
| 12. | Office of Naval Research (Code 482)<br>Naval Ocean Research & Development<br>Activity<br>NSTL Station<br>Bay St. Louis, MS 39529           | 1 |
| 13. | Scientific Liaison Office<br>Office of Naval Research<br>Scripps Institution of Oceanography<br>La Jolla, CA 92037                         | 1 |
| 14. | Library<br>Scripps Institution of Oceanography<br>P.O. Box 2367<br>La Jolla, CA 92037  | 1 |
| 15. | Library<br>Department of Oceanography<br>University of Washington<br>Seattle, WA 98105   | 1 |
| 16. | Library<br>CICESE<br>P.O. Box 4803<br>San Ysidro, CA 92073   | 1 |
| 17. | Library<br>School of Oceanography<br>Oregon State University<br>Corvallis, OR 97331  | 1 |
| 18. | Commander<br>Oceanographic Systems Pacific<br>Box 1390<br>Pearl Harbor, HI 96860   | 1 |
| 19. | Chief, Ocean Services Division<br>National Oceanic and Atmospheric<br>Administration<br>8060 Thirteenth Street<br>Silver Springs, MD 20910 | 1 |

- |     |  |   |
|-----|--|---|
| 20. | Chairman, Oceanography Department<br>U.S. Naval Academy<br>Annapolis, MD 21402   | 1 |
| 21. | Director<br>Naval Oceanography Division (OP952)<br>Navy Department<br>Washington, D.C., 20350                          | 1 |
| 22. | Mr. Ben Cagle<br>Office of Naval Research Branch Office<br>1030 East Green Street<br>Pasadena, CA 91106                | 1 |
| 23. | Dr. Robert E. Stevenson<br>Scientific Liaison Office, ONR<br>Scripps Institution of Oceanography<br>La Jolla, CA 92037 | 1 |
| 24. | Mr. Jerry Norton, Code 68<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, CA 93940             | 1 |
| 25. | Ms. Bonita Hunter, Code 68<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, CA 93940            | 1 |
| 26. | Mr. Laurence Breaker<br>National Environmental Satellite Service<br>660 Price Avenue<br>Redwood City, CA 94063         | 1 |
| 27. | LT. Carol Jori, Code 68<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, CA 93940               | 1 |
| 28. | LT. Walter Hanson<br>USCG Oceanographic Unit, Bldg. 159-E<br>Navy Yard Annex<br>Washington, D.C. 20593                 | 1 |



DATE  
FILMED  
— 8